


4. ALGEMENE VOËLS / GENERAL BIRDS


[WAARGENOME VOËLSPECIES IN DIE KOUE BOKKEVELD, WARM BOKKEVELD EN TANKWA-KAROO]
[OBSERVED BIRD SPECIES IN THE KOUE BOKKEVELD, WARM BOKKEVELD AND TANQUA-KAROO]

4a. Algemeen, medium-grootte / General, medium-sized

4b. Algemeen, medium: Rivier-en vleiland / General, medium: River and wetland



Columba livia | Tuinduif | Rock Dove
348 | 33 | WB | S | LC PD



Columba guinea | Kramduif | Speckled Pigeon
190 | 13 | KB | WB | TK | S | LC PS



Columba arquatrix | Geelbekbosduif | African Olive-Pigeon
349 | 37-42 | KB | WB | S | LC PD



Streptopelia semitorquata | Grootringduif | Red-eyed Dove
352 | 30-33 | KB | WB | S | LC PI



Streptopelia capicola | Tortelduif (Gewone) | Cape Turtle-Dove
354 | 25-28 | KB | WB | TK | S | VU



Oena capensis | Namakwaduijie | Namaqua Dove
356 | 28 | KB | WB | TK | S | LC PI



Aplopelia larvata | Kaneelduif | Lemon Dove
360 | 24-26 | SM | S | LC PS




Streptopelia senegalensis | Rooiborsduif | Laughing Dove
355 | 22-24 | KB | WB | TK | S | LC PS



Chrysococcyx klaas | Meitjie | Klaas's Cuckoo
383 | 16-18 | KB | WB | TK | S | LC PS



Chrysococcyx caprius | Diederikkie | Diederik Cuckoo
386 | 17-19 | KB | WB | TK | M | LC PS



Centropus burchellii | Gewone vleioerie | Burchell's Cuckoo
391 | 40-41 | SM | S | LC PS




Centropus viridis | Gewone vleioerie | Burchell's Cuckoo
391 | 40-41 | SM | S | LC PS



Collus striatus | Gevlekte muisvoël | Speckled Mousebird
424 | 13 | WB | S | LC PI




Collus collus | Witrusmuisvoël | White-backed Mousebird
425 | 13 | KB | WB | TK | S | LC PI



Urocolinus indicus | Rooiwangmuisvoël | Red-faced Mousebird
426 | 34 | KB | WB | TK | S | LC PS



Mirafra apicalis | Kaapse Klapperjie | European Bee-eater
438 | 25-28 | KB | WB | TK | M | LC PS



Upupa africana | Hoephoep | African Hoopoe
451 | 25-28 | KB | WB | TK | M | LC PD



Tricholerna leucomelas | Bonthoekapper | Acacia Pied Barbet
465 | 16-18 | WB | TK | S | LC PD



Geocolaptes olivaceus | Grondspieg | Ground Woodpecker
480 | 25 | WB | TK | S | NT



Dendropicos ariseoscephalus | Gryskoppag | Olive Woodpecker
488 | 16-18 | SM | S | LC PS



Mirafra apicalis | Kaapse Klapperjie | Cape Clapper Lark
495 | 19 | KB | WB | TK | S | LC PD



Dicurus adsimilis | Mikstertbyvanger | Fork-tailed Drongo
541 | 22-26 | KB | SM | S | LC PS



Pycnonotus capensis | Kaapse Tiptol | Cape Bulbul
566 | 19-21 | KB | WB | TK | S | LC PS



Pycnonotus nigricans | Rooioogtiptol | African Red-eyed Bulbul
567 | 19-21 | TK | S | LC PI



Cossypha caffra | Gewone Janfrederik | Cape Robin-Chat
601 | 16-17 | KB | WB | TK | S | LC PS



Lanius ferrugineus | Suidelike waterfiskaal | Southern Boubon
736 | 20-22 | WB | S | LC PS



Corcomela familiaris | Gewone Spekrevter | Familiar Chat
589 | 14-15 | KB | WB | TK | S | LC PS



Corcomela frater | Woeslynspakrevter | Frater Chat
590 | 14-15 | TK | S | LC PS



Corcomela sinuata | Vlaaktepskrevter | Sickle-winged Chat
591 | 14-15 | KB | TK | S | LC PS



Corcomela schlegelii | Karospekrevter | Karoo Chat
592 | 16-18 | WB | TK | S | LC PS



Sigelus silens | Fiskaalvlievanger | Fiscal Flycatcher
698 | 17-20 | WB | TK | S | LC PS



Lanius collaris | Fiskaalklaksman (Gewone) | Common Fiscal
732 | 21-23 | KB | WB | TK | S | LC PI



Onychognathus nabevanger | Bleekvlervanger | Pale-winged Starling
770 | 26-28 | KB | TK | S | LC PS



Tetoporus zeylonus | Bokmakierie | Bokmakierie
746 | 22-23 | KB | WB | TK | S | LC PS



Sturnus vulgaris | Europese Spreu | Common Starling
757 | 20-22 | KB | WB | TK | S | LC PD



Spreo bicolor | Witgatspreeu | Pied Starling
758 | 27-28 | KB | WB | TK | S | LC PS



Creatophora cinerea | Lepspreu | Wattled Starling
760 | 19-21 | KB | WB | TK | S | LC PI



Onychognathus morio | Rooivlekvanger | Red-winged Starling
769 | 27-30 | KB | WB | TK | S | LC PI



Onychognathus morio | Rooivlekvanger | Red-winged Starling
769 | 27-30 | KB | WB | TK | S | LC PI



Alcedo semitorquata | Blouvisvanger | Half-collared Kingfisher
430 | 18-20 | SM | S | LC PD




Tchagra tchagra | Grysborsstjagra | Southern Tchagra
742 | 17-21 | SM | S | LC PI



Quelea quelea | Rooibekwelea | Red-billed Quelea
821 | 11-13 | TK | N | LC PS



Emberiza impetiana | Vaalstreepkoppe | Lark-like Bunting
887 | 74 | KB | WB | TK | S | LC PS



Ceryle rudis | Bontvisvanger | Pied Kingfisher
428 | 23-25 | KB | WB | TK | S | LC PS



Megascops asio | Reusevisvanger | Giant Kingfisher
429 | 38-43 | KB | WB | S | LC PD



Alcedo semitorquata | Blouvisvanger | Half-collared Kingfisher
430 | 18-20 | SM | S | LC PD



Alcedo semitorquata | Blouvisvanger | Half-collared Kingfisher
430 | 18-20 | SM | S | LC PD



Alcedo cristata | Kufkopvisvanger | Malachite Kingfisher
431 | 13-14 | KB | WB | TK | S | LC PS




Halcyon albinervis | Bruinkopvisvanger | Brown-hooded Kingfisher
435 | 19-20 | SM | S | LC PS




Motacilla capensis | Gewone Kwikkie | Cape Wagtail
713 | 19-20 | KB | WB | TK | S | LC PS



Ploceus capensis | Kaapse Wewer | Cape Weaver
813 | 17 | KB | WB | TK | S | LC PS



Ploceus velatus | Swarkeelgeelvink | Southern Masked Weaver
814 | 17-18 | KB | WB | TK | S | LC PS




Euplectes orix | Rooivink | Southern Red Bishop
824 | 10-11 | KB | WB | TK | S | LC PS



Euplectes capensis | Kaaapse Flap | Yellow Bishop
827 | 12 | KB | WB | TK | S | LC PS



Indicator indicator | Groothuigsvanger | Greater Honeyguide
474 | 13-22 | WB | TK | S | LC PS



Indicator minor | Kleinhuigsvanger | Lesser Honeyguide
476 | 13-15 | TK | S | LC PS



Andropadus impatorius | Gewone Willie | Sombre Greenbul
572 | 15-18 | SM | S | LC PS

VERKLARING/KEY

Wetenskaplike naam/ Scientific name | STANDSTATUS (RESIDENSIE/RESIDENCE)
Afrikaanse gebruiknaam | N= Nomadies/Vagrant
M= Migrerend/Migrant
S= Standvoël/Resident bird | Common English name

Podiceps cristatus | Kuifkopdobber | Great Crested Grebe
6 | 45-56 | KB | WB | TK | SM | S | LC PD

Roberts VII no | LENGTE/Length (cm) | WAAKSTADUS/STATUS (CONSERVATION STATUS)
CR = Critically endangered
EN = Endangered
VU = Vulnerable
NT = Near threatened
LC = Least concern
LC PI = Least concern: Population increasing
LC PD = Least concern: Population declining

STREEK (Area)/REGION (Area) | WARM BOKKEVELD/Warm Bokkeveld
KB: Koue Bokkeveld | WB: Warm Bokkeveld
TK: Tanqua Karoo
SM: Skarpsig gebied/Rare migrant

BRONNE/SOURCES

Sinclair I, Hockey P, Tarboton W 2002. *SASOL Voëls van Suider-Afrika*. 3 Ed. Struik, Kaapstad.
Hockey P, Dean WRJ, Ryan 2005. *Roberts birds of Southern Africa*. 7 Ed. Percy FitzPatrick Institute of African Ornithology, University of Cape Town.
Andreanidis A: *Scientific Names of Southern African Birds Explained: Is it all Greek to you?*
International Union for the Conservation of Nature (IUCN)
BEELDE/IMAGERY
Encyclopedia of Life (eol.org)
grobimages.co.za
http://www.biodiversityexplorer.org/birds/
https://en.wikipedia.org/wiki/trevorhardaker.co.za
warwicktarboton.co.za
www.arkive.org

